

BACTERINA POLIVALENTE

DESCRIPTION

BACTERINA POLIVALENTE is an inactivated bacterine in injectable suspension containing *Actinobacillus lignerisesi*, *Corynebacterium pyogenes*, *Staphylococcus aureus*, *Salmonella abortus* and *Pasteurella multocida*.

TECHNICAL PROPERTIES

All pathological diseases of reproductive, lymphatic and mastitic character are pathologically related to three microorganisms: *Actinobacillus lignerisesi*, *Corynebacterium pyogenes* and *Staphylococcus aureus*. It is common to find polyvalent infections caused by the combined action of some of them.

Salmonella abortus is a germ that causes paratífic abortus in sheep and goat. *Pasteurella multocida* is the causative agent of pneumonia and mastitis in sheep and goats.

INDICATIONS AND SPECIES

Intended for sheep, goats and cattle of all ages for the prevention of:

- Actinobacillosis for sheep and cattle
- Sheep mastitis, pyobacillar mastitis and cattle staphylococia.
- Paratífic abortion in sheep and cattle.
- Enteritis and colibacilosis in lambs, calves and young goats.
- Pneumoenteritis.

COMPOSITION

| | |
|-----------------------------------|------------|
| <i>Actinobacillus lignerisesi</i> | 3000 m.CFU |
| <i>Staphylococcus aureus</i> | 3000 m.CFU |
| <i>Corynebacterium pyogenes</i> | 3000 m.CFU |
| <i>E. coli</i> | 3000 m.CFU |
| <i>Salmonella abortus</i> | 3000 m.CFU |
| <i>Pasteurella multocida</i> | 3000 m.CFU |
| Excipient to 1 ml | |

DOSAGE

It is administered, after shaking the vial, intramuscularly at the following doses:

- Lambs: 0.5 ml at two days of age. Revaccinate with 1 ml two weeks later and with 1.5 ml at 15 klw (kilogram of live weight)
- Calf: 2 ml at two days of age. Revaccinate with 2 ml two weeks later and with 10 ml at 150 klw

C/Luxemburg, 25 – Polígono
Industrial
Apdo 15 . 43120 CONSTANTÍ
(Tarragona)
Tel. 977 29 63 04 Fax 977 29 63 05
www.biovét-alquermes.com

Premio a la Internacionalización
Cámara de Comercio 2005



INTERNATIONAL ADDRESS
P.O Box 15
43120 CONSTANTÍ (Tarragona)
SPAIN
Tel. 34 – 977 29 63 04
Fax 34 – 977 29 63 05

- Sheep for reposition: 2 ml two months before the first heat and re-vaccinate with 2 ml three weeks before the first mating.
- Cattle: 4 ml two months before the first heat and revaccinate with 2 ml three weeks before the first mating.
- Reproductive animals: Sheep (2 ml); and cattle (4 ml) three weeks before and three weeks after parturition. Males every 6 months.

The vaccination schedule must be established under veterinary criteria.

CONTRAINDICATIONS

Do not administer to sick animals, with fever or submitted to immuno suppressing treatments.

PACKAGING

250 ml glass vials. MUST BE MAINTAINED BETWEEN 4 AND 8 °C.

WITHDRAWAL PERIOD

Meat: 21 days.

C/Luxemburg, 25 – Polígono
Industrial
Apdo 15 . 43120 CONSTANTÍ
(Tarragona)
Tel. 977 29 63 04 Fax 977 29 63 05
www.biovet-alquermes.com

Premio a la Internacionalización
Cámara de Comercio 2005



INTERNATIONAL ADDRESS
P.O Box 15
43120 CONSTANTÍ (Tarragona)
SPAIN
Tel. 34 – 977 29 63 04
Fax 34 – 977 29 63 05