

NEOBACTERINA PORCINA REPRO

DESCRIPTION

NEOBACTERINA PORCINA REPRO is an inactivated bacterine in injectable suspension containing *Actinobacillus suis*, *E.coli*, *Corynebacterium pyogenes*, *Salmonella enteritidis* and *Pasteurella multocida*.

TECHNICAL PROPERTIES

With increasing intensity, pigs present pathological processes with diarrhea, pneumonia, dermatitis and nephritis. These diseases have been grouped in two classes: respiratory diseases and dermatitis-nephropathy syndrome. Both share a common bacterial origin: *Actinobacillus suis*.

Actinobacillus suis produces cardiac, renal and lung hemorrhagic lesions, as well as lymphatic lesions (immunosuppression).

E.coli, *Corynebacterium pyogenes* and *Salmonella enteritidis* are secondary germs associated with most infectious reproductive processes (miscarriage, embryonic mortality...) and with digestive processes of early ages.

Pasteurella multocida is a secondary germ associated with respiratory diseases.

INDICATIONS AND SPECIES

Intended for pigs for the prevention of:

- Digestive problems: enteritis and salmonellosis
- Reproductive problems: metritis, miscarriage, infectious metritis
- Respiratory diseases (pasteurelosis) in young animals
- Dermatitis-nephropathy syndrome
- Improvement of the immune system of the animal.

COMPOSITION

<i>Actinobacillus suis</i>	2000 m. CFU
<i>Actinobacillus pleuroneumoniae</i>	1000 m. CFU
<i>Bordetella bronchiseptica</i>	3000 m. CFU
<i>Pasteurella haemolytica</i>	3000 m. CFU
<i>Salmonella enteritidis</i>	3000 m. CFU
Excipient to	1 ml

DOSAGE

It is administered, after shaking the vial, with a subcutaneous injection at the following dose: piglets, 0.5 ml to 1 ml; fattening and reproductive, 2 ml.

C/Luxemburg, 25 – Polígono Industrial
Apdo 15 . 43120 CONSTANTÍ (Tarragona)
Tel. 977 29 63 04 Fax 977 29 63 05
www.bioviet-alquermes.com

Premio a la Internacionalización
Cámara de Comercio 2005



INTERNATIONAL ADDRESS
P.O Box 15
43120 CONSTANTÍ (Tarragona)
SPAIN
Tel. 34 – 977 29 63 04
Fax 34 – 977 29 63 05

The recommended vaccination schedule in farms with reproductive and digestive problems is the following:

	Piglets from non-vaccinated mothers	Piglets from vaccinated mothers		Breeders	Males
1 st or 2 nd day of life	0.5 ml	-	5 months	2 ml	-
14 days	1 ml	1 ml	Rest	2 ml 2 weeks before labour and 2 ml 3 weeks after it	2 ml twice a year
Before weaning	1 ml	1 ml			

CONTRAINDICATIONS

Do not administer to animals that are sick, have fever or are subjected to immuno suppressing treatments.

PACKAGING

250 ml glass vials. MUST BE MAINTAINED BETWEEN 4 AND 8 °C.

WITHDRAWAL PERIOD

Meat: 21 days.

C/Luxemburg, 25 – Polígono Industrial
Apdo 15 . 43120 CONSTANTÍ (Tarragona)
Tel. 977 29 63 04 Fax 977 29 63 05
www.bioviet-alquermes.com

Premio a la Internacionalización
Cámara de Comercio 2005



INTERNATIONAL ADDRESS
P.O Box 15
43120 CONSTANTÍ (Tarragona)
SPAIN
Tel. 34 – 977 29 63 04
Fax 34 – 977 29 63 05