

NEOBACTERINA PORCINA NEUMO

DESCRIPTION

NEOBACTERINA PORCINA NEUMO is an inactivated bacterine in injectable suspension containing *Actinobacillus suis*, *Actinobacillus (haemophilus) pleuroneumoniae*, *Bordetella bronchiseptica*, *Pasteurella haemolytica* and *Salmonella enteritidis*.

TECHNICAL PROPERTIES

With increasing intensity, pigs present pathological processes with diarrhoea, pneumonia, dermatitis and nephritis. These diseases have been grouped in two classes: Respiratory diseases and dermatitis-nephropathy syndrome both with a common bacterial origin: *Actinobacillus suis*.

Actinobacillus suis produces cardiac, renal and lung haemorrhagic lesions, as well as lymphatic lesions (immunosuppression).

Actinobacillus (haemophilus) pleuroneumoniae produces pleuroneumonia and can be found in articular lesions and dermatitis.

Bordetella bronchiseptica, *Pasteurella haemolytica* and *Salmonella enteritidis* are secondary germs associated to most infections and respiratory processes.

INDICATIONS AND SPECIES

Intended for pigs for the prevention of piglet pneumoenteritis, atrophic rhinitis, respiratory diseases: pasteurellosis and pneumonias, dermatitis-nephropathy syndrome and for the improvement of the immune system functions.

COMPOSITION

<i>Actinobacillus suis</i>	2000 m. CFU
<i>Actinobacillus pleuroneumoniae</i>	1000 m. CFU
<i>Bordetella bronchiseptica</i>	3000 m. CFU
<i>Pasteurella haemolytica</i>	3000 m. CFU
<i>Salmonella enteritidis</i>	3000 m. CFU
Excipient to	1 ml

DOSAGE

It is administered, after shaking the vial, subcutaneously at the following doses: piglets, 0.5 ml to 1 ml; fattening and reproductive animals, 2 ml.

C/Luxemburg, 25 – Polígono
Industrial
Apdo 15 . 43120 CONSTANTÍ
(Tarragona)
Tel. 977 29 63 04 Fax 977 29 63 05
www.bioviet-alquermes.com

Premio a la Internacionalización
Cámara de Comercio 2005



INTERNATIONAL ADDRESS
P.O Box 15
43120 CONSTANTÍ (Tarragona)
SPAIN
Tel. 34 – 977 29 63 04
Fax 34 – 977 29 63 05

The recommended vaccination schedule in farms with respiratory problems is the following:

	Piglets from non-vaccinated mothers	Piglets from vaccinated mothers		Fattening	Males
1 st or 2 nd day of life	0.5 ml	-	5 months	2 ml	-
14 days	1 ml	1 ml	Rest	2 ml 2 weeks before labour and 2 ml 3 weeks after it	2 ml twice a year
Before feed	1 ml	1 ml			

CONTRAINDICATIONS

Do not administer to animals that are sick, have fever or are subjected to immuno suppressing treatments

PACKAGING

250 ml glass vials. MUST BE MAINTAINED BETWEEN 4 AND 8 °C.

WITHDRAWAL PERIOD

Meat: 21 days.

C/Luxemburg, 25 – Polígono Industrial
 Apdo 15 . 43120 CONSTANTÍ (Tarragona)
 Tel. 977 29 63 04 Fax 977 29 63 05
 www.biovet-alquermes.com

Premio a la Internacionalización
 Cámara de Comercio 2005



INTERNATIONAL ADDRESS
 P.O Box 15
 43120 CONSTANTÍ (Tarragona)
 SPAIN
 Tel. 34 – 977 29 63 04
 Fax 34 – 977 29 63 05