

## SULVET 48

### DESCRIPTION

Sulphamethoxyipyridacine and trimethoprim in oral and injectable solution.

### TECHNICAL PROPERTIES

The association of sulphamides and trimethoprim is widely known because of the synergism that exists between them. It possesses a wide action spectrum for accompanying germs of mycoplasmosis: *Staphylococcus*, *Streptococcus* and *Pasteurella*. It has a good organic diffusion.

Sulphamethoxyipyridacine is a sulphamide with a good absorption and distribution, achieving high pulmonary concentrations. It acts through competitive antagonism with para-aminobenzoic acid.

### INDICATIONS AND SPECIES

Intended for poultry, pigs, sheep, goats and cattle in the treatment of :

- Respiratory diseases: C.R.D of poultry, rhinitis, pharyngitis, sinusitis, bronchopneumonia, acute and chronic bronchitis, pasteurelosis.
- Gastrointestinal diseases: enteritis, peritonitis, pneumoenteric diseases and gastritis.
- Local and general infections.

### COMPOSITION

Sodium sulphamethoxyipyridacine	40 mg
Trimethoprim	8 mg
Excipient to	1 ml

### DOSAGE

DEEP INTRAMUSCULAR injection: at doses of 1 ml/ 20 k.l.w. every 24 hours for 4- 5 days

ORALLY: 0.5 ml per litre in drinking water.

### INCOMPATIBILITIES

Do not administer together with streptomycin, tetracyclines, kanamycin or with rehydrating solutions containing lactates or chlorides. It increases the plasma concentration of teophelines and the anticoagulant effect of warfarines

### CONTRAINDICATIONS

Do not administer to animals with hepatic or renal problems or to animals allergic to sulphamides. Prolonged treatment can affect the folic acid metabolism, therefore it is recommended to administer vitamin B complex.

C/Luxemburg, 25 – Polígono  
Industrial  
Apdo 15 . 43120 CONSTANTÍ  
(Tarragona)  
Tel. 977 29 63 04 Fax 977 29 63 05  
www.bioviet-alquermes.com

Premio a la Internacionalización  
Cámara de Comercio 2005



INTERNATIONAL ADDRESS  
P.O Box 15  
43120 CONSTANTÍ (Tarragona)  
SPAIN  
Tel. 34 – 977 29 63 04  
Fax 34 – 977 29 63 05

### **OVERDOSAGE**

It can produce neuritis or renal lesions to ruminants which should be treated with alkalinizers of the urine.

### **PACKAGING**

Injectable solution: 100 and 250 ml  
Oral solution: 1 and 5 litres.

### **WITHDRAWAL PERIOD**

Meat: 5 days.  
Milk: 3 days.

C/Luxemburg, 25 – Polígono  
Industrial  
Apdo 15 . 43120 CONSTANTÍ  
(Tarragona)  
Tel. 977 29 63 04 Fax 977 29 63 05  
[www.biovét-alquermes.com](http://www.biovét-alquermes.com)

Premio a la Internacionalización  
Cámara de Comercio 2005



**INTERNATIONAL ADDRESS**  
P.O Box 15  
43120 CONSTANTÍ (Tarragona)  
SPAIN  
Tel. 34 – 977 29 63 04  
Fax 34 – 977 29 63 05